

**Effect of Using Organic Substance Safe (Rigir) and Artificial Feeding
on Growth Performance of *Oreochromis niloticus* and Grey Mullet
(*Mugil cephalus*).**

Ahmed F. Fath El-Bab

Central Laboratory for Aquaculture Research at Abbassa, Agriculture Research
Center.

ABSTRACT

This study was carried out to investigate the effect of organic fertilizer with treated poultry manure as safe organic matter (rigirs) at rate of 100kg/feddian weekly and artificial feed on growth performance and economic efficiency as well as pond productivity of Nile tilapia (*Oreochromis niloticus*) and grey mullet (*Mugil cephalus*). Ten ponds (21×100 m-½ feddan) were stocked by Nile tilapia and grey mullet. Ten ponds (two replicates for each treatment) were designed for five treatments and stocked with *Oreochromis niloticus* (4000 fish/pond), grey mullet (2000 fish/pond). The experimental earthen ponds used in the study were treatment 1(control): artificial feeding throughout the experimental period (26 weeks), treatment 2: ponds fertilized by (rigirs) at rate of 50kg/ pond weekly for the first month of the experimental period, followed by pelleted artificial feed until the end of experiment, treatment 3: ponds fertilized by (rigirs) at rate of 50kg/ pond weekly for the first two months of the experimental period, followed by pelleted artificial feed until the end of experiment, treatment 4: ponds fertilized by (rigirs) at rate of 50kg/ pond weekly for the first three month of the experimental period, followed by artificial feed until end of experiment, and treatment 5: ponds fertilized by (rigirs) at rate of 50kg/ pond weekly during the whole experimental period (26 weeks). Results obtained are summarized in the following: Growth parameters: Nile tilapia (ON): Treatment 1 recorded the highest ranges of body weight (BW), body length (BL), daily weight gain (DWG) and specific growth rate (SGR), while treatment 3 recorded the highest ranges of condition factors (K). Grey mullet (GM): Treatment 1 recorded the highest ranges of body weight (BW), body length (BL), daily weight gain (DWG) and specific growth rate (SGR), while treatment 2 recorded the highest ranges of condition factors (K). Chemical composition: Nile tilapia: treatment 1 recorded the highest ranges of fat and ash%, treatment 4 recorded the highest ranges of protein content. While, treatment 2 recorded the highest ranges of moisture. Grey mullet: treatment 4 recorded the highest ranges of moisture and protein content. treatment 1 recorded the highest fat content, while treatment 2 recorded the highest ranges of ash%. Economical evaluation, content of the obtained results showed that, the use of rigir at the first three months of culture Nile tilapia and grey mullet at rate of 100 kg / feddan / week, after that use artificial feed containing 25% crude protein at rate of 3 % of total biomass of in earthen ponds could be recommended for producing Nile tilapia and grey mullet. From the economical point of view, this treatment seemed to be the best in terms of ratio of returns to total costs.

Keywords: Nile Tilapia - Grey Mullet - Rigir

INTRODUCTION

Oreochromis niloticus and grey mullet (*Mugil cephalus*) are considered in

Egypt as fish species of high market value. Afifi *et al.* (1996) reported that both tilapia and mullet responded in their growth

FATH EL-BAB

performance when they stocked together in earthen ponds fertilized with chicken manure, super phosphate and urea with supplementary diet containing 11% crude protein. Ease cultivation of tilapia and mullet, resistance to poor water quality and disease, tolerance to a wide range of environmental conditions, ability to convert efficiently organic domestic and agricultural wastes into high quality protein, good growth rates and amenability to intensification are some of the basic characteristics of both species which make them ideal candidates for intensive and semi-intensive culture (Afifi *et al.*, 1996).

Grey mullet is commonly cultured with tilapia in Egypt. Tilapia and mullet are popular and favored by the Egyptian consumers for their good quality flesh and comparatively larger size of their adults. Both species are highly priced and feasible culture in fishponds. One way of increasing the food availability in fishponds is fertilization of ponds by chemical or/and organic fertilizers which is often a mean of increasing the primary natural productivity of the ponds as reported by FAO (1980). The objective of the present study aimed to evaluate the effect of rigir without artificial feeding, and comparing with artificial feeding on the production of Nile tilapia and Grey mullet fingerlings in the intermediate fish farming for youth project.

MATERIALS AND METHODS

The study had been done in a private farm (in Tollumbat No. 7 in Riyad City, Kafr El-Sheikh governorate, Delta district at the Northern part of Egypt) to evaluate the effect of fertilization and artificial feeding on earthen ponds productivity stocked with Nile tilapia (*Oreochromis niloticus*) and grey mullet (*Mugil cephalus*). Rigirs is a product

for fish farm produced by Misr El-Salam International Company for producing organic fertilizer, Alexandria governorate. Rigirs consists of chicken manure but compressed and heat treated in order to be free from parasites, Salmonella, Shigella and *E. coli*. The procedures done in this study such as fertilizers application per treatment, pond preparation, stocking rate and pond daily management are described in details. Also water quality measurements, fish sampling and data collected during harvest are recorded too. Equations and statistical methods for analysing the specific growth rate, daily weight gain and the condition factor are given. The current experiment was conducted using randomized block design for five treatments of similar surface area (2100 m²) each

Ponds description and preparation

The experimental ponds were equal in water volume (2625 m³) and dimensions (21x100 m) with the same average water depth of 125 cm. Before beginning the experiment, ponds were drained, weeded and exposed to the sun for one month for complete dryness. Each pond had inlet and outlet water gates through which the water level was controlled.

The farm water source was mainly agricultural drainage water and comes from El-Gharbia drainage canal. The water system of the experimental ponds is maintained by gravity. Ponds inlet and outlet pipes were covered with narrow mesh screen to prevent unwanted fish or predators to get into ponds. Partial filling of the ponds to 50% of target level started on the following day after applying the initial fertilization dose from relevant fertilizers. Two days prior to stocking

ORGANIC SUBSTANCE AND ARTIFICIAL FEED IN NILE TILAPIA AND GREY MULLET PRODUCTION

fingerlings fish, ponds water level increased and reached the maximum target water depth.

Experimental fish

The experimental ponds were stocked with fish species; *Oreochromus niloticus* and *Mugil cephalus*. *O. niloticus* fingerlings were stocked at an average initial total length of 9.44cm and an average initial total weight of 10.20g for all treatments. The average initial total length of G. mullet fingerlings were 13.94cm and an average initial total weight of 13.80g for all treatments. The fingerlings of *O. niloticus* and *M.cephalus* were collected from different fish farms Riyad City, Kafer El-Shiek Governorate. Each pond was stocked with 6000 fish (4000 *O. niloticus* and 2000 *M.cephalus*) corresponding to the rate of 12000 fish/ feddan. The trial lasted for 182 days started on the 15th of April and harvested on 14th October 2013.

Fertilizers applications

Ponds were fertilized for the twenty sixth weeks. Fertilization occurred once a week by broadcasting of rigir. The chemical composition of rigir is show in Table (1).

Organic fertilizer and Supplementary feed

Rigirs is a product for fish farm produced by Misr El-Salam International Company for producing oraganic fertilizer, Alexandria Governorate. Rigirs consists of chicken manure but compressed and heat treated in order to be free from parasites, Salmonella, Shigella and *E. coli*. at rate of 50kg/ (pond-½feddan) /weekly.

Commercial diet was manufactured by Sherbeen-Domiatte, local factory. Sample of fish feed was collected from several sacks and send for proximate analysis at the Central Laboratory for Aquaculture Research at Abbassa. Chemical analysis of experimental

artificial feed is presented in Table (2). The fingerlings were fed commercial floating diet to keep the diets available for fish contain 25% crude protein (pellets 3 mm in diameter), and fed six days per week at a daily feeding rate of 3% of the estimated fish-weight twice at 9.00am and 3.00pm during the experimental period. Feed was applied by broadcasting over pond water surface in the same place and fish were considered satiated when they did not show an interest on the feed.

Table 1. Chemical analysis of rigirs on D.M basis.

Item	Rigir
Cubic meter weight (kg/ m3)	730
Humidity %	9.6
pH	8.01
Electric conductivity mmhos/cm	4.2
Total nitrogen %	2.38
Ammonia nitrogen ppm	1040
Organic matter %	59.68
Organic carbon %	29.58
Ash %	40.32
C:N ratio	13:1
Total phosphorus %	1.79
Total potassium %	1.91
Iron ppm	768
Manganese ppm	398
Copper ppm	40
Parasites	Nil

FATH EL-BAB

Table 2: Composition and proximate chemical analysis of the experimental ration.

Ingredients	%
Yellow corn	33
Wheat bran	17
Fish meal (72% CP)	10
Meat meal	7
Soybean meal (44% CP)	10
Decorticated cotton seed meal	7
Poultry slaughter by-products	8
Fat	5
Vitamin premix*	1.5
Mineral premix**	1.5
Total	100
Proximate analysis (% of dry weight)	
Moisture	10.20
Crude protein	22.08
Ether extract	6.54
Crude fiber	6.72
Ash	6.06

* Each gram of vitamin premix contains 20000 IU vit. A , 2000 IU vit.D₃, 400 IU vit E, 20 mg Niacin, 4.5 mg riboflavin, 3 mg pyridoxine, 0.013 mg vit. B12, 100mg choline chloride and 2mg vit K.

** each gram contains 0.83 mg Ca, 0.63 mg P, 0.78 mg Na, 0.018 mg Zn and 0.001 mg Cu. The Mixture was prepared by mixing 35 parts of dicalcium phosphate, 3 parts of mineral premix and 2 parts of common salt

Feed quantity was adjusted according to average body weight of the sample in each pond. In order to determine the average weight of fish, biweekly samples were taken by seining, where 30 fishes / species from each pond (replicate) were collected and then retained again to the pond after individual measuring the weight and length.

Water management

Water temperature, dissolved oxygen and pH were measured daily at 6 a.m. and 12 p.m. using thermometer, dissolved oxygen

meter (YSI model 57) and pH meter (model Corning 345), respectively. Determinations of the other water quality parameters (alkalinity, nitrate, nitrite and ammonia) were carried out every two weeks according to the methods of Boyd (1979).

Fish samples and measurements

Random samples (30 fish from each species for each pond -120 fish for each treatment) were taken every 15 days during the experimental period. During this experiment, body measurements (body weight in g and body length in cm) at biweekly interval throughout the whole experiment period.

Condition factor was determined by using the following formula:

$$K = \frac{\text{weight (g)}}{\text{length (cm)}^3} \times 100$$

Specific growth rate was calculated according to Jauncey and Rose (1982) by using the following formula:

$$SGR = \frac{\ln W_2 - \ln W_1}{t} \times 100$$

Harvesting

At the end of the experiment (14th of October, 2013), ponds were gradually drained from the water and fish were harvested by seining and transferred to fiberglass tanks and carried to the processing centre where they washed, and the fish of the different fish species were sorted and collectively weighed.

Chemical analysis of fish

At the end of the experiment, five fish were chosen from each species from each pond and exposed to the chemical

ORGANIC SUBSTANCE AND ARTIFICIAL FEED IN NILE TILAPIA AND GREY MULLET PRODUCTION

composition of whole fish body according to the methods of A O A C (1990).

Statistical analysis

The statistical analysis of data collected was carried out by applying the computer program (SAS, 1996). Differences among means were tested for significance according to Duncan's multiple range test (1955).

RESULTS AND DISCUSSION

Water quality parameters

Results of water quality parameters of the experimental ponds during the experimental period (182 days) as averages of the monthly samples are summarized in Table (3). In general, averages of water temperature ranged from 26.48 to 27.63°C during the experiment course (15th April – 15th October). Gui *et al.* (1989) and Kamal *et al.* (2008)

found that an average temperature of 28 °C was optimal for growth of Nile tilapia fry.

Dissolved oxygen ranged between 4.21 and 6.89 mg / l. Denzer (1968), AIT (1986), and Hassan *et al.* (1997) reported that 2.3 mg DO / l is above the normal tolerance level of tilapia.

The value of pH ranged between 8.12 and 8.80. Boyd (1998) reported that waters with a pH range of 6.5 – 9 are the most suitable for fish production.

The average concentration of unionized ammonia (NH₃) was 0.35, 0.32, 0.29, 0.28 and 0.26 mg/l for T1, T2, T3, T4 and T5, respectively. Diana and Lin (1998) reported ammonia concentration of 0.374 – 0.410 mg/l in ponds fertilized with both chicken manure and inorganic fertilizers in combination

Table 3: Some water quality parameters of earthen ponds stocked with Nile tilapia and Grey mullet fry treated with organic fertilizers and different types of diets for 182 days culture period.

Variable	NO.	T1	T2	T3	T4	T5
Temperature (C ^o)	6	27.63±0.50a	27.18±0.50a	26.48±0.50b	27.13±0.50a	26.52±0.50b
DO oxygen	6	5.48±0.15ab	6.89±0.15a	5.50±0.15ab	5.42±0.15ab	4.21±0.15b
PH	6	8.80±0.12a	8.56±0.12a	8.52±0.12ab	8.37±0.12b	8.12±0.12b
Seechi disk (cm)	6	28.28±0.15a	26.19±0.15a	25.06±0.15ab	24.52±0.15b	20.39±0.15b
NH ₃ mg/l	6	0.35±0.008b	0.32±0.008b	0.29±0.008a	0.28±0.008a	0.26±0.008a
NO ₂ mg/l	6	0.033±0.001c	0.040±0.001a	0.036±0.001b	0.038±0.00a	0.035±0.001b
NO ₃ mg/l	6	0.112±0.003a	0.110±0.003a	0.117±0.003a	0.115±0.003a	0.115±0.003a
Total alkalinity (mg/l)	6	422.97±14.06a	413.55±14.06a	402.63±14.06ab	392.96±14.06ab	384.76±14.06b

± Means with the same letter in each column are not significantly different ($P \geq 0.05$)

FATH EL-BAB

The concentration of nitrite ranged between (0.033 to 0.040 mg/l) during the whole experimental period. The concentration of nitrate seems to be constant among treatments (0.110 to 0.117 mg/l). Kamal *et al.* (2008) found that, the values of nitrite ranged between 0.02 and 0.03 mg/L. Diana and Lin (1998) noticed that nitrite and nitrate concentration ranging between 0.374 – 0.410 mg/l and 0.438 – 0.461 mg/l, respectively, in ponds fertilized with both chicken manure and inorganic fertilizers.

The average value of seechi disk readings were 18.28, 19.16, 22.06, 21.52 and 23.39 cm for T1, T2, T3, T4 and T5, respectively. The values of the total alkalinity ranged between 384.76 and 422.97 mg/l, and total hardness ranged between 191.16 to 212.36 mg/l. Kamal *et al.* (2008) reported that, the values of the total alkalinity ranged between 392.50 and 434.34 mg/l, and. The above results showed that all parameters of water quality were in the suitable range (Boyd, 1979).

Body weight

The averages of initial weights of *Oreochromis niloticus* (ON) and Grey mullet *Mugil cephalus* (GM) were 47.11 and 36.87g respectively; while at the end of the experiment, the means of body weight for fish species were 307.90 and 23.45g, respectively (Table 4). These results indicate that, the body weight for (ON) was higher than obtained in (GM).

The initial body weight was 47.10, 47.95, 46.33, 45.89 and 48.30g for the different treatments T1 (control): artificial feeding during the experimental period, T2: fertilized with rigirs for only the first month of

the experimental period, followed by pelleted artificial feed until the end of experiment, T3: fertilized with rigirs for the first two months of the experimental period, followed by pelleted artificial feed until the end of experiment, T4: fertilized by rigirs for the first three monthes of the experimental period, followed by artificial feed until end of experiment, and T5: fertilized with rigirs till the end of the experimental period, respectively. While at the end of experiment, the means of tilapia body weight for treatments were 366.20, 351.40, 338.77, 288.80 and 194.40g for the five treatments, respectively. These results indicate that, the average body weight for first and second treatment was higher than other treatments. Osman *et al.* (2008) found that, fish groups fed the artificial diet only (treatment 1) showed the highest significant finale body weights ($P<0.01$) followed in descending order by treatments 3, 4, 5 and 2 (308.8, 292.4, 234.1, 202.3 and 178.8 g), respectively.

Green (1992) concluded that the chicken manure can replace 100% of pelleted supplemental feed without significant effects on the growth of tilapia during the first 60 days of the culture. This is in agreement with the results obtained in the present study. Fath El-bab *et al.* (2011) studied the effect of fertilization and artificial feed on productivity of earthen ponds stoked under polyculture system and found that, the differences between fish species were significant. These results may be attributed to the feeding habits of species as tilapia fish is an efficient converter of phytoplankton and can utilize a wide variety of food, especially artificial feeds.

Concerning body weight of grey mullet (Table 4), the initial body weight was

ORGANIC SUBSTANCE AND ARTIFICIAL FEED IN NILE TILAPIA AND GREY MULLET PRODUCTION

36.31, 38.12, 37.24, 35.14 and 37.55g for the different treatments, respectively. While at the end of experiment, the means of body weight for treatments were 278.68, 268.13, 256.57, 227.40 and 146.45g for the five treatments, respectively. These results indicate that, the average body weight for first treatment was higher than other treatments and the differences among treatments were significant ($P<0.05$).

El -Ebiary (1998) when studied the use of organic manures in polyculture system for tilapia, mullet and carp, he found that, the average of final weight influenced by an increase of the period of organic fertilization. He referred that, organic fertilizer is not enough to rely upon as a staple food for Fish.

Jasmin *et al.* (2011) found that, average gain in body weight of all the fish species together was less in the pond without fertilizer than that in the pond with fertilizer. The fertilizer increased the primary productivity and finally caused a significant increase in fish yield in ponds.

Body length

Table 5 shows means of total body length, at the start and the end of experiment. As described in this Table the initial averages of ON and GM were 16.87 and 17.22cm, respectively while at the end of experiment, the means of total body length for fish species were 26.94 and 29.98cm, respectively. These results indicate that body length for GM was significantly higher than ON.

Concerning the effect of treatments on body length for Nile tilapia (Table 5), the initial body length was 16.00, 17.80, 15.80, 16.50 and 18.20cm for five treatments T1, T2, T3, T4 and T5 respectively. While at the end of the experiment, the means of body length for treatments were 28.10, 27.90, 27.40, 26.10 and 25.20cm for five treatments, respectively. These results indicate that the total body length for first treatment (control) was higher than other treatments. Analysis of variance of results indicated that the differences among treatments were significant ($P<0.05$).

Table 4: Least-square means and standard error of the tested factors affecting on body weight (gm).

Treatments	No.	Nile tilapia		Grey mullet	
		Initial weight	Final weight	Initial weight	Final weight
T1	30	10.02±0.92a	366.20±1.29a	16.50±0.92b	318.68±1.29a
T2	30	10.60±0.92 a	351.40±1.29ab	17.20±0.92a	308.13±1.29a
T3	30	10.49±0.92a	338.77±1.29b	17.09±0.92a	296.57±1.29ab
T4	30	10.01±0.92 a	288.80±1.29c	16.61±0.92b	288.40±1.29b
T5	30	9.90±0.92 b	194.40±1.29d	16.67±0.92b	156.45±1.29c
Overall mean		10.20±1.04	307.90±2.58	16.80±1.04	273.65±2.58

± Means with the same letter in each column are not significantly different ($P\geq 0.05$)

FATH EL-BAB

Table 5: Least-square means and standard errors of the tested factors affecting total body length (cm).

Treatments	No.	Nile tilapia		Grey mullet	
		Initial length	Final length	Initial length	Final length
T1	30	9.30±0.09a	28.10±0.78a	12.82±0.09b	32.30±0.78a
T2	30	9.72±0.09a	27.90±0.78a	13.22±0.09a	31.70±0.78b
T3	30	9.69±0.09a	27.40±0.78b	13.19±0.09a	31.64±0.78b
T4	30	9.34±0.09a	26.10±0.78c	12.80±0.09b	31.86±0.78b
T5	30	9.18±0.09a	25.20±0.78d	12.68±0.09b	27.55±0.78c
Overall mean		9.44±0.41	26.94±1.35	12.94±0.41	30.99±1.35

± Means with the same letter in each column are not significantly different ($P \geq 0.05$)

Osman *et al*, (2008) reported that, average fish final length at the end of the experimental period was found to follow the same order ($P < 0.01$) of final body weight and being 31.20, 30.80, 30.50, 29.95 and 27.38cm, respectively. These results are in accordance with those obtained by Hafez (1991), who found a strong correlation between body weight and body length for tilapia, mullet and carp fish.

Averages mullet body length at the start of this experiment were 17.10, 17.70, 18.00, 16.80 and 16.50cm for five treatments, respectively. While at the end of experiment the means of body length for treatments were 31.30, 30.70, 31.60, 30.80 and 26.50cm for the five treatments, respectively. These results indicate that, the average body length for first treatment was higher than other treatments. Prabahar and Murugan (2012) found increasing body length when mullet and major carp, fed on artificial feed after 2 months of fertilization by organic and inorganic fertilizers.

Condition factor (K)

Condition factor (K) was considered to be a sufficient measure of shape, although shape is usually not considered as a character of interest to breeding programs, since it has no obvious economic value (Nilsson, 1992).

Table 6 presents means of K at the start and the end of the experiment as affected by treatments and fish species. As described in this Table, the averages of K values at the start for ON and GM were 1.17 and 0.74, respectively while at the end of experiment the means of K for fish species were 1.83 and 1.02, respectively.

With regard to the effect of treatments on condition factor for Nile tilapia, Table 6 showed that, the initial condition factor was 1.43, 1.00, 1.47, 1.04 and 0.94 for five treatments T1, T2, T3, T4 and T5 respectively. While at the experimental end, the averages of condition factor for treatments were 1.97, 1.95, 2.00, 1.89 and 1.61, respectively. These results indicate that, the

ORGANIC SUBSTANCE AND ARTIFICIAL FEED IN NILE TILAPIA AND GREY MULLET PRODUCTION

Table 6: Least-square means and standard error of the tested factors affecting on condition factor (K).

Treatments	No.	Nile tilapia		Grey mullet	
		Initial K	Final K	Initial K	Final K
T1	30	1.24±0.04a	1.65±0.18a	0.79±0.04a	0.95±0.18a
T2	30	1.15±0.04b	1.62±0.18a	0.74±0.04a	0.97±0.18a
T3	30	1.15±0.04b	1.69±0.18a	0.74±0.04a	0.94±0.18a
T4	30	1.22±0.04a	1.58±0.18a	0.79±0.04a	0.90±0.18b
T5	30	1.28±0.04a	1.21±0.18b	0.81±0.04a	0.75±0.18c
Overall mean		1.21±0.11	1.57±0.49	0.78±0.11	0.92±0.49

± Means with the same letter in each column are not significantly different ($P \geq 0.05$).

condition factor for T3 was higher than other treatments. Analysis of variance of results indicates that, the differences among treatments were significant ($P < 0.05$). Oren (1981) revealed that fluctuations in ponds for fish growth (length, weight and condition factor) are affected by different factors such as feeding regime, population density and environmental conditions. Osman *et al.* (2008) when studied the effect of different combinations between organic, inorganic fertilizers and artificial diets on the growth performance, feed efficiency and production cost of *Oreochromis niloticus* reared in earthen ponds, found that the average of condition factor values for treatments (Treatment1, artificial diet only), (Treatment2, fertilization only), (Treatment3, 1month fertilization + 3 months artificial diet), (Treatment4, 2months fertilization + 2 months artificial diet) and (Treatment5, 3 months fertilization + 1 month artificial diet) were 2.38, 2.34, 2.4, 2.26 and 2.18, respectively.

Results presented in Table 6 show that, the average of condition factor values for treatments (1, 2, 3, 4 and 5) For grey

mullet at the beginning of this experiment were 0.75, 0.70, 0.66, 0.77 and 0.81, respectively. While at the of this experimental period the average of condition factor values were 1.05, 1.07, 1.04, 1.00 and 0.95 for five treatments, respectively, the differences among treatments were significant ($P < 0.05$).

Fath El-bab *et al.* (2011) found that, feeding treatment in adequate quantities and the increase in feeding rate resulted in higher condition factor since the fish grow well when the supply of food is adequate. Similar results in which condition factors increased with the feeding rate have been reported by Chua and Teng (1982).

Daily weight gain (DWG)

Table 7 shows means of daily weight gain, throughout the experiment as affected by treatments and fish species. As described in this Table the averages between each of ON and GM were 1.43 and 1.11g/fish, respectively. These results indicate the daily weight gain for ON was higher than obtained for GM.

FATH EL-BAB

Table 7: Least-square means and standard error of the tested factors affecting on DWG and SGR.

Treatments	No.	Nile tilapia		Grey mullet	
		DWG, g/fish	SGR, %/d	DWG, g/fish	SGR, %/day
T1	30	1.96±0.07a	2.00±0.12a	1.67±0.07a	1.73±0.12a
T2	30	1.87±0.07ab	1.92±0.12ab	1.63±0.07a	1.68±0.12ab
T3	30	1.80±0.07ab	1.85±0.12b	1.53±0.07b	1.61±0.12b
T4	30	1.53±0.07c	1.57±0.12c	1.49±0.07bc	1.57±0.12c
T5	30	1.01±0.07d	1.05±0.12d	0.77±0.07c	0.84±0.12d
Overall mean		1.63±0.30	1.68±0.20	1.41±0.30	1.48±0.20

± Means with the same letter in each column are not significantly different ($P \geq 0.05$).

Concerning the effect of treatments on daily weight gain for Nile tilapia Table 6 showed that, the daily weight gain was 1.75, 1.67, 1.61, 1.33 and 0.80g/fish for five treatments T1, T2, T3, T4 and T5, respectively. These results indicate that, the daily weight gain for treatments T1 and T2 were higher than other treatments. Analysis of variance of results indicates that the differences among treatments were significant ($P < 0.05$). Osman *et al.* (2008) found that, the highest averages daily gain and SGR were recorded by treatment 1 (2.54 and 3.62%/ day, respectively) followed by treatments 3,4 and 5, while the lowest values (1.46 and 3.16%/day, respectively) were found in the group fed no artificial feed (treatment 2).

Hassan *et al.* (2008) revealed that artificial feed application to blue tilapia reared in earthen ponds improved specific growth rate values as compared with other treatments in fertilized ponds at lower stocking densities. Increasing was recorded in the growth performance by using supplemental feeding compared with other

treatments (fertilized ponds). They found also that, blue tilapia was the fastest growing fish under these treatment conditions. On the other hand, growth performance of grass carp was higher in fertilized ponds.

With regard to the effect of treatments on daily weight gain for grey mullet, it was 1.33, 1.26, 1.20, 1.17 and 0.61,g/fish for five treatments, respectively. These results indicate that, the daily weight gain for treatments T1 and T2 were higher than other treatments. Analysis of variance of results indicates that the differences among treatments were significant ($P < 0.05$). El - Ebiary (1998) when studied the use of organic manures in polyculture system for tilapia, mullet and carp, he found that the average of daily weight gain Influenced by an increase of the period of organic fertilization. He referred that, organic fertilizer is not enough to rely upon as a staple food for Fish. Kstmont (1995) reported that, the growth and yield of each species may be high in polyculture than in monoculture because of positive interactions among species.

ORGANIC SUBSTANCE AND ARTIFICIAL FEED IN NILE TILAPIA AND GREY MULLET PRODUCTION

Specific growth rate (SGR)

Table 7 shows means of SGR, at the end of experiment as affected by treatments and fish specie. As described in this Table, the averages of ON and GM were 1.68 and 1.48%/day, respectively. These results indicate that SGR for ON was higher than that obtained for GM.

With regard to the effect of treatments on SGR for Nile tilapia, Table 7 showed that, the specific growth rate during the whole experimental period were 1.13, 1.10, 1.09, 1.00 and 0.77 for five treatments T1, T2, T3, T4 and T5, respectively. These results indicate that, the Specific growth rate for T1 and T2 were higher than those obtained with other treatments. Analysis of variance indicates that the differences among treatments were significant ($P < 0.05$). Abdel-Rahman *et al* (2003) reported that SGR for *O. niloticus* during the fortnight periods of the experiment were significantly higher for pelleted and mash treatments rather than chicken manure and cow manure treatments.

Hassan *et al.* (2008) reported that, the SGR values of blue tilapia influenced by supplementary feed which increased the availability of all nutrients to fish when stocked in un-fertilized ponds; the lowest value was obtained by grass carp with high stocking density in un-fertilized ponds.

The average of SGR values for treatments (1, 2, 3, 4 and 5) For grey mullet during the whole experimental period were 1.12, 1.07, 1.06, 0.97 and 0.78, respectively. These results indicate that, the SGR for treatments T1 was higher than other treatments. Analysis of variance of results indicates that the differences among treatments were significant ($P < 0.05$). Sarig (1981) observed that mass development of

algae frequently appears in the fertilized ponds where supplementary food is added. He added that these algae are not fully utilized by common carp or tilapia, but serve as excellent food for grey mullet. Essa *et al.* (1989) and Hussein (1995) reported that, the highest growth of mullet (*M. cephalus*) was recorded in polyculture conditions with organic fertilizers and supplementary feeding.

Chemical composition

Moisture %

Table 8 showed averages of moisture% for whole body at the end of this experiment found to be 67.31 and 63.90% for ON and GM, respectively. These results indicated that, the moisture% for ON was higher than that obtained for GM. The differences between fish species were significant ($P < 0.05$).

With regard to the effect of treatments on moisture% for Nile tilapia, Table 8 showed that, the values at the end of this experiment were 67.18, 69.36, 67.75, 66.22 and 66.04% for five treatments T1, T2, T3, T4 and T5, respectively. These results indicate the dry matter for T2 was higher than obtained for other treatments. Analysis of variance of results indicates that the differences among treatments were significant ($P < 0.05$). Osman *et al.* (2008) reported that, moisture content of tilapia fish in treatment2 (only organic and inorganic fertilizers) decreased significantly ($P < 0.01$) compared with the other treatments.

Concerning the effect of treatments on moisture% for grey mullet, Table 8 showed also that, the moisture% was 64.31, 64.25, 62.44, 64.73 and 63.79%/fish for five treatments T1, T2, T3, T4 and T5 respectively. These results indicate that, the moisture% for treatments 4 was higher than

FATH EL-BAB

Table 8: Least-square means and standard error of the tested factors affecting chemical composition.

Treatments	No.	Moisture	Protein	E. ext.	Ash
Nile tilapia					
T1	6	67.18±1.29b	59.37±1.05d	31.75±0.79a	12.68±0.64a
T2	6	69.36±1.29a	65.54±1.05b	29.53±0.79c	11.12±0.64ab
T3	6	67.75±1.29b	63.33±1.05c	30.95±0.79b	11.47±0.64ab
T4	6	66.22±1.29c	66.08±1.05a	29.25±0.79c	11.03±0.64ab
T5	6	66.04±1.29c	66.73±1.05a	27.52±0.79d	10.95±0.64b
Overall mean		67.31±1.41	64.21±1.11	29.80±0.88	11.45±0.74
Grey mullet					
T1	6	64.31±1.68a	64.92±1.38c	32.52±1.02a	12.69±0.83b
T2	6	64.25±1.68a	64.97±1.38c	28.85±1.02b	13.60±0.83a
T3	6	62.44±1.68c	65.29±1.38b	28.69±1.02b	13.48±0.83a
T4	6	64.73±1.68a	65.99±1.38b	26.75±1.02c	11.37±0.83c
T5	6	63.79±1.68b	66.22±1.38a	22.69±1.02d	11.16±0.83d
Overall mean		63.90±1.71	65.48±1.40	27.95±1.05	12.66±0.84

± Means with the same letter in each column are not significantly different ($P \geq 0.05$).

other treatments. Analysis of variance of results indicates that the differences among treatments were significant ($P < 0.05$).

Protein %

The averages of protein content of ON and GM were found to be 63.21 and 64.48%, respectively. These results indicate that, the protein % for GM was higher than obtained in ON. The differences between fish species were significant ($P < 0.05$).

With regard to the effect of treatments for Nile tilapia on protein %, Table 8 showed that, protein content was found to be 58.37, 64.54, 62.33, 65.08 and 65.73% for five treatments T1, T2, T3, T4 and T5, respectively. These results indicated that, protein % for T5 was higher than obtained with other treatments. Analysis of variance of results indicates that the differences among treatments were significant ($P < 0.05$).

Hafez *et al.* (2001) found that, fish flesh from fish fed the first feeding treatment

(organic fertilization) had higher percentages of protein, ash and lower percentages of fat compared with flesh obtained from the second feeding treatment (supplementary feeds).

The average of protein % values for treatments (1, 2, 3, 4 and 5) For grey mullet at the end of the experimental period were 63.92, 63.97, 64.29, 64.99 and 65.22%, respectively. These results indicate that, the % protein for treatments T5 was higher than other treatments. Analysis of variance of results indicates that the differences among treatments were significant ($P < 0.05$).

Ether extract %

As described in Table (8), the averages fat content of ON and GM were 26.80 and 24.95%, respectively. These results indicate the ether extract for ON was higher than that obtained for Gm and the differences between fish species were significant ($P < 0.05$).

ORGANIC SUBSTANCE AND ARTIFICIAL FEED IN NILE TILAPIA AND GREY MULLET PRODUCTION

With regard to the effect of treatments for Nile tilapia on ether extract % Table 8 showed that, ether extract at the end of this experiment was 28.75, 26.53, 27.95, 26.25 and 24.52% for five treatments T1, T2, T3, T4 and T5, respectively. These results indicate that ether extract for first treatment was higher than obtained for other treatments, and by increasing fat percentage there was decreasing protein percentage. Analysis of variance indicates that the differences between treatments were significant ($P>0.05$). Barash and Schroeder (1984) indicated that tilapia fat content was higher in fish fed pelleted feed when compared by those raised with fermented manure

The average of ether extract% values for treatments (1, 2, 3, 4 and 5) For grey mullet during the whole experimental period was found to be 29.52, 25.85, 25.69, 23.75 and 19.69%, respectively. These results indicate that, the ether extract% for first treatments was higher than other treatments. Analysis of variance of results indicates that the differences among treatments were significant ($P<0.05$).

Ash %

The averages Ash content of ON and GM were found to be 10.45 and 11.66%, respectively. These results indicated that the ash for GM was higher than obtained for ON. The differences among fish species were significant ($P<0.05$).

With regard to the effect of treatments for Nile tilapia on ash%, Table 8 showed that ash content was found to be 11.68, 10.12, 10.47, 10.03 and 9.95% for treatments T1, T2, T3, T4 and T5, respectively. These results indicate that ash for first treatment (T1) was higher than obtained for other treatments, and the differences among

treatments were significant ($P>0.05$). Brown and Murphy (1991) concluded that larger size fish class (as treatment 1 in the present experiment) usually had lower ash and higher fat content than smaller size (treatment 2).

The average of ash% values for treatments (1, 2, 3, 4 and 5) For grey mullet during the whole experimental period were 11.69, 12.60, 12.48, 10.37 and 10.16%, respectively. These results indicate that, ash% for second treatment (T2) was higher than other treatments. Analysis of variance of results indicates that the differences among treatments were significant ($P<0.05$). It is obvious to notice that the rest in fish body composition was high in groups treated with fertilizers, this may be due to the increase of natural food in fish gut.

Total yield

Averages of total yield at the end of the experiment were listed in Table (9). As described in this Table, *Oreochromis niloticus* gained the highest yield 8806.93kg - 100% compared with 2234.55kg - 25.37% gained by grey mullet. These results may be attributed to the feeding habits of the two species as described previously. The total fish production (tilapia fish + grey mullet) for all feeding treatments was 12693.22 kg. Fath El-Bab *et al.* (2011) found that, this increase may be due to the dietary habits of each species.

As in this Table, Nile tilapia (T1) gained the highest yield (2219.46kg), compared with (T2) (2117.63kg), (T3) (2030.74kg), (T4) (1486.94kg) and (T5) (952.16kg). While total yield of grey mullet, (T1) gained the highest yield (540.64 kg), compared with (T2) (509.45kg), (T3) (487.48kg), (T4) (427.51kg) and (T5) (269.47kg) These results indicated that, the

FATH EL-BAB

Table 9: Total yield of different fish species as affected by feeding treatments.

Treatments	Nile tilapia		Grey mullet	
	Yield (kg)	%total yield	Yield (kg)	%total yield
T1	2219.46	25.20%	958.61	24.71%
T2	2117.63	24.04%	947.32	24.42%
T3	2030.74	23.06%	913.14	23.54%
T4	1486.94	16.88%	875.69	22.57%
T5	952.16	10.81%	729.22	18.79%
Overall mean	8806.93	100%	3879.84	100
% of the biggest value	100%		44.05	

**The fish catch attributed to the highest value for the same variable.*

low of gain in rigid ponds compared with control (fed diet) reported in similar studies (Green, 1992, Diana *et al.*, 1994; 1996 and Brown *et al.*, 2000) indicates that either phytoplankton may not be enough to meet protein requirement of fish or that fish could not efficiently assimilate the produced phytoplankton in these ponds. Collis and Smitherman (1978) found that hybrid tilapia when fed on manure, grew 62% compared to hybrids fed on a high protein diet. Barash and Schroeder (1984) found that the substitution of 46% of the pellets by fermented cow manure did not reduce the total fish yield but the complete substitution of the pellets by fermented cow manure caused 47% decrease in the total yield.

Economic evaluation

Results of costs including variable, fixed and interest on working capital for the 196

treatments applied are shown in Table (10). Results of this Table revealed that costs of fish fingerlings are similar in all treatments applied, however, the feed costs differed according to rigid with some diets and thus, the net return was the lowest for treatment 5, rigid (1124.85LE) and increased to 5554.45, 5177.92, 5633.33 and 7259.05 LE for other diets treatment 1, treatment 2, treatment 3 and treatment 4, respectively.

Total costs per feddan increased from 15799.48LE (100%) for treatment 5 to 43258.59 (274%), 40955.08 (259%), 37316.56 (236%) and 24926.94 (158%), respectively. Differences in total costs were attributed to the differences in feed costs and rigid.

Total return in LE/feddan for treatment 1, treatment 2, treatment 3, treatment 4 and treatment 5 were 49371.93, 46691.9, 43508.78, 32744.88 and 16924.33 LE, respectively (Table 10). Net

ORGANIC SUBSTANCE AND ARTIFICIAL FEED IN NILE TILAPIA AND GREY MULLET PRODUCTION

Table 10: The effect of the experimental diets on economic efficiency (LE/Feddan)

Items	Treatments					
	T1	T2	T3	T3	T5	
1- Variable costs (LE/½Feddan)						
a. costs of fish fingerlings:						
O. niloticus	1600	1600	1600	1600	1600	
G. Mullet	5520	5520	5520	5520	5520	
b. Feeds	30618.2	28475.0	25089.7	13562.3	0	
c. rigir	0	780	1560	2340	5070	
Total variable costs (LE/Feddan)	37738.24	35595.03	32209.72	20682.29	12190	
2- Fixed costs (LE/Feddan)						
a. Depreciation (materials&others) 10%	300	300	300	300	300	
b. Taxes	250	250	250	250	250	
Total fixed costs (LE/Feddan)	550	550	550	550	550	
Total operating costs (variable&fixed)	38288.24	36145.03	32759.72	21232.29	12740	
Interest on working capital *	2863.751	2703.45	2450.247	1588.059	952.8822	
Total costs	41151.99	38848.48	35209.97	22820.35	13692.88	
% of the smallest value	300	284	257	167	100	
Returns						
Total return (LE) **	ON	25745.69	24246.89	23840.86	16133.34	8037.93
	GM	23626.24	22445.01	19667.93	16611.54	7352.63
	Total	49371.93	46691.9	43508.78	32744.88	15390.55
Net return (LE/Feddan)	8219.94	7843.41	8298.82	9924.54	1697.67	
% of the smallest value of net return	917.2166	875.2026	926.0186	1107.423	100	
% Net returns to total costs	19.97	20.19	23.57	43.49	6.54	

* $15\% \times \text{total operating costs} \times 140/365 \text{ days}$.

** *The economical evaluation of results was carried out according to market prices in 2012 in LE.*

O. niloticus = LE 160 /1000 fry.

G. mullet = LE 920 /1000 fry

Rigir = LE 1950/1000 Kg

Fish feed (25% protein) = LE 3200 /1000 Kg

returns / feddan in LE were found to be. 5554.45, 5177.92, 5633.33, 7259.05 and 1124.85LE for the five treatments T1, T2, T3, T4 and T5, respectively. The percentage of net return to total costs 14.13, 14.07, 16.59, 31.36 and 7.12% for treatments T1, T2, T3, T4 and T5, respectively.

These results indicated that feeding of *Oreochromis niloticus* in polyculture with grey mullet in earthen ponds on natural feed (rigir) for 3 months followed by feeding on an artificial feed containing 25% crude protein resulted in best economic efficiency compared to the other treatments. These results are in

FATH EL-BAB

complete agreement with results of Kamal *et al.* (2008).

CONCLUSION

Based on results obtained in this study and on the economical evaluation, it could be concluded that, the use of rigir at the first three months of culture tilapia and grey mullet at rate of 100 kg / feddan / week (as safe organic matter), after that use artificial feed containing 25% crude protein at rate of 3 % of total biomass of Nile tilapia and grey mullet in earthen fish ponds culture at last three months could be recommended for producing Nile tilapia and grey mullet. From the economical point of view, this treatment seemed to be the best in terms of ratio of returns to total costs..

REFERENCES

- Abdel-Rahman, A. A., M. M. Abdel- Aal, Fatma A. Hafez; Sohair A. Arafa and G. M. Abdel-Aziz. (2003).** Effect of commercial diets, manure and some agriculture by- products on performance of *Oreochromis niloticus* in polyculture system. Egyptian Soc Anim. Reprod. Fert. Fifteenth Annual Congr. 26–30, January, 2003, pp. 195- 208.
- Affi, E. A., Fatma A. Hafez, Abdel-Hakim, N. F. and Abdella, M. M. (1996).** Effect of mullet stocking rate on some productive traits in fish. Egypt. J. Agric. Res., 74(2):2127-2143.
- (AIT) Asian Institute of Technology (1986).** Buffalo/ fish and duck/fish integrated systems for small – scale farmers at the family level. AIT Research Report No. 198. Bangkok, Thailand, 138 pp.
- AOAC (1990).** Official Methods of Analysis. 15th Ed. Published by the Association of Official Analytical Chemists. North Nineteen st. Auite 210 Arlingth, Virginia 2220/USA.
- Barash, J. E. and G. L. Schroeder. (1984).** Use of fermented cow manure as substrate for fish polyculture in stagnate water ponds. Aquaculture, 36: 127-140.
- Boyd, C. E. (1979).** Water Quality in Warm Water Fish Ponds. Ed Claude E. Boyd. Third printing, 1984. Pub. Auburn Univ., Agri.Exp. Station, AID/Dsan- G.G.00 39.pp. 359.
- Boyed, C. E. (1998).** Water quality for pond aquaculture. Research and development series No. 43. pp. 37. International Centre for aquaculture and aquatic Environments. Alabama Agricultural Experiment Station. Auburn University.
- Brown, M. L. and B. R. Murphy. (1991).** Relationship of relative weight (Wr) to proximate composition of juvenile striped bass and hybrid striped bass. Trans. Am. Fish. Soc., 120: 509-518.
- Brown, C. L., R. B. Bolivar, E. T. Jimenez and J. Szype. (2000).** Timing of the onset of supplemental feeding of Nile tilapia (*Oreochromis niloticus*) in ponds. Page 237 in K. Fitzsimmons, editor. Proceedings of the Fifth International Symposium on Tilapia Aquaculture. 3-7 September, Rio De Janeiro. Brazil.
- Chua, T. E. and S. K. Teng. 1982.** Effects of food ration on growth, condition factor, food conversion efficiency, and net yield of estuary grouper, *Epinephelus salmonides* Maxwell, cultured in floating Net-cages. Aquaculture, 27: 273-283.
- Collis, W.J. and R.O. Smitherman. (1978).** Production of tilapia hybrids with cattle manure of a commercial diet. pp 43-54. In R. O. Smitherman, W. L. Shelton and J. H. Grover (editors). Culture of exotic fishes symposium proceedings. Fish culture section, American Fisheries Society, Auburn, Alabama.
- Denser, H. W. (1968).** Studies on the physiology of young tilapia. FAO Fisheries Report, 44 (4): 357 – 366.

ORGANIC SUBSTANCE AND ARTIFICIAL FEED IN NILE TILAPIA AND GREY MULLET PRODUCTION

- Diana, J. S., C. K. Lin and K. Jaiyen (1994).** Supplemental Feeding of tilapia in fertilized ponds. *J. world Aquaculture. Soc.* 25 (4): 497-506.
- Diana, J. S., C. K. Lin and Y. Yi (1996).** Timing of Supplemental Feeding for tilapia in production. *J. world Aquaculture Soc.*, 27, (4): 410-419.
- Diana, J. S., and C. K. Lin (1998).** the effects of fertilization and water management on growth and production of Nile tilapia in deep ponds during the dry season. *J. of the World Aquaculture Society.* 29 (4) 405 – 413.
- Duncan, D.B. (1955).** Multiple range and multiple F test. *Biometrics*, 11:1-42.
- Essa, M.A; El-Sherif, Z.M.; Abu El-Ezz, S.A. and Abd El-Moati, A.R. (1989).** Effect of water quality, food and viability and crowding on rearing conditions and growth parameter of some economical fish species grown under polyculture systems. *Bull. Nat. Inst. Ocean. & Fish. ARE*, 15(1): 125-134,
- FAO (1980).** Food and Agriculture Organization of the United Nation). *Fish Feed Technology.* ADCP/PER/80/11. Aquaculture development and co-ordination program.
- Fath El-bab A. F; Samra I. M.; Mostafa M. A. A. and Gouda A. H. (2011).** The effect of fertilization and artificial feeding on productivity of earthen ponds stocked under polyculture system. *Abbassa Int. J. Aqua.*, Vol.5 No.(1): 500-527.
- Green, B. W. 1992.** Substitution of organic manure for pelleted feed in tilapia production. *Aquaculture*, 101:213-222.
- Gui,Y., Z. Wang, Y. Chen, W. Zheng and F. Li. (1989).** Use of fluctuating temperature to promote growth of *Tilapia nilotica*. *Journal of Fisheries of China.* 13 (4) 326 – 331.
- Hafez, A. Fatma (1991).** Studies on some Productive Traits in Fish. pp.63-85. Ph.D. Thesis, Faculty of Agriculture at Moshtohor. Zagazig University, Egypt.
- Hafez, A. Fatma, M. A. Soltan and M.K. Ibrahim. (2001).** Effect of organic fertilization, supplementary feeding and stoking rate on carcass and chemical composition of Nile tilapia and silver carp. *Egyptian Journal Agricultural Resours*, 78 (4): 2128-2143.
- Hassan, S. Amal; S.H. Sayed and E. M. Ibrahim. (2008).** Effects of stoking and supplementary feed on the growth performance of plue tilapia (*Oreochromis aureus*) and grass carp (*Ctenoharyngodon idella*) reared in earthen ponds.
- Hassan, S., P. Edwards and D. C. Little. (1997).** Comparison of tilapia monoculture and carp polyculture in fertilized earthen ponds. *J. Worled Aquaculture Society*, 28 (3): 268 – 274.
- Hussein, M.S. (1995)** Semi-intensive fish culture in some Egyptian fish farms, ph. D. Thesis. Fac. Agr. Suez Canal Univ, 256 pp.
- Jasmin, S.; F. Ahmed; S. H. Rahman; M. A. S. Jewel and M. Y. Hossain. (2011).** Effect of organic and inorganic fertilizers on growth performance of carps in earthen ponds through polyculture system. *Our Nature.* 9: 16-20.
- Jauncey, K. and B. Rose. (1982).** A guide to tilapia feeds and feeding. Institute of Aquaculture, University of Sterling, Scotland.
- Kamal, S. M; Abdel Ghanny, S. and Abdel All, M. M. (2008):** Effect of nutrition and fertilization on production of Nile Tilapia (*O. niloticus*) fingerlings in concrete ponds. Proceedings of the 8th international symposium on Tilapia in aquaculture, Volume (1), pp: 387-401. Cairo International Convention Center (CICC), Egypt, 12-14 October, 2008.
- Kamal, S. M., A. A. Mahmoud and R. A. Abou-Seif (2009):** Effect of organic fertilizer and different supplementary feeds on production of Nile tilapia (*O. niloticus* L.) and Silver carp (*Hypophthalmichthys molitrix* val.) in earthen ponds. *Abbassa International Journal for*

FATH EL-BAB

- Aquaculture. Special issue for global fisheries and aquaculture research conference. Cairo International Convention Center (CICC), Egypt, 24-26 October, 2009 pp: 570-586.
- Kstmont, P. 1995.** Different systems of carp production and their impacts on the environment. *Aquaculture*, 129: 347-372.
- Nilsson, J., (1992):** Genetic parameters of growth and sexual maturity in Arctic char (*Salvelinus alpinus*). *Aquacultur*, 106:9-19.
- Oren, O. H. (1981).** Aquaculture of grey mullets. International. Biological Programme. Cambridge Univ. London. Pp.45-57.
- Osman, M. F., Khattab, H. M., Mounes, H. A. M. And Hafez, F. A. (2008):** Productive performance of *Oreochromis niloticus* under different nutritional and aquaculture systems. International Symposium on Tilapia in Aquaculture, 887-902.
- Prabaha, C. and Senthil Murugan. (2012).** A study on effects of fertilization and supplementary feed on growth performance of major carps. *International Journal of Pharmaceutical & Biology Archives*. 3(1): 186-191.
- Ramdan A. A., M. A. El- Nady, M. G. Kamar and F. A. Hafez. (2001).** Effect of different husbandry methods on water quality and growth of some fresh water fishes. *Egypt. J. Aquat. Boil. & Fish*. 5 (4):295-318.
- Sarig, S. (1981)** The Mugilidae in polyculture in fresh and brackish water fishponds. 391 - 393 in *Aquaculture of grey mullets* . Edited by O.H. Oren, Israel Ocean. And Limnol Res., 507 pp.
- SAS (1996).** SAS procedure Guide "Version 6.12 ed". SAS Institute Inc., Cray, NC, USA.

ORGANIC SUBSTANCE AND ARTIFICIAL FEED IN NILE TILAPIA AND GREY MULLET
PRODUCTION

تأثير استخدام مادة عضوية آمنة (ريجر) والتغذية الصناعية على أداء النمو في البلطي النيلي
والبورى

أحمد فاروق فتح الباب

المعمل المركزى لبحوث الثروة السمكية ، مركز البحوث الزراعية ، وزارة الزراعة ، مصر.

تم إجراء هذه الدراسة لمعرفة تأثير التسميد العضوي بزرق الدواجن المعامل حراريا ومكبوس (ريجر) ١٠٠ كجم/ اسبوعيا/ الفدان والأعلاف الصناعية على أداء النمو والكفاءة الاقتصادية لأسماك البلطي والبورى في أحواض ترابية فى نظام الاستزراع السمكى متعدد النوع، تم تخزين الأسماك فى عشرة أحواض (٢١ × ١٠٠ م) أسماك البلطي النيلي والبورى. تم تصميم العشرة أحواض (مكررين لكل معاملة) لتشمل خمسة معاملات وتم تخزين أسماك البلطي النيلي (٤٠٠٠ سمكة / حوض) مع أسماك البورى (٢٠٠٠ سمكة / حوض)، وكانت الأحواض الترابية التجريبية التي استخدمت في الدراسة المعاملة الاولى (الكنترول): التغذية الصناعية طوال فترة التجربة (٢٦ أسبوعا)، المعاملة الثانية: تم التسميد بواسطة ريجر بمعدل ٥٠ كجم / الأسبوع/ الحوض طوال الشهر الأول من بداية التجربة، تليها التغذية الصناعية حتى نهاية فترة التجربة، المعاملة الثالثة: تم التسميد بواسطة ريجر بمعدل ٥٠ كجم / الأسبوع/ الحوض لمدة شهرين من بداية التجربة، تليها التغذية الصناعية حتى نهاية فترة التجربة، المعاملة الرابعة: تم التسميد بواسطة ريجر بمعدل ٥٠ كجم / الأسبوع/ الحوض طوال ثلاثة شهور من بداية التجربة، تليها التغذية الصناعية حتى نهاية فترة التجربة و المعاملة الخامسة: تم التسميد بواسطة ريجر بمعدل ٥٠ كجم / لأسبوع/ الحوض طوال فترة التجربة (٢٦ أسبوعا). ويمكن تلخيص النتائج المتحصل عليها فيما يلي:

قياسات النمو:

أسماك البلطي: سجلت المعاملة الاولى أعلى معدل من وزن الجسم الحى، طول الجسم، زيادة اليومية فى الوزن و معدل النمو النوعى وسجلت المعاملة الثالثة أعلى معدل لمعامل الحالة،

أسماك البورى: سجلت المعاملة الاولى أعلى معدل من وزن الجسم الحى، طول الجسم، زيادة اليومية فى الوزن و معدل النمو النوعى وسجلت المعاملة الثانية أعلى معدل لمعامل الحالة،

FATH EL-BAB

التركيب الكيميائي:

أسماك البلطي: سجلت المعاملة الاولى أعلى معدل لنسبة الرطوبة والدهن وسجلت المعاملة الخامسة أعلى معدل لنسبة الروتين، و نسبة الرماد .

أسماك البورى: سجلت المعاملة الاولى أعلى معدل لنسبة الدهن وسجلت المعاملة الثالثة أعلى معدل لنسب الرطوبة وسجلت المعاملة الرابعة أعلى معدل لنسب الرماد وسجلت المعاملة الخامسة أعلى معدل لنسبة الروتين

إستنادا إلى النتائج التي تم الحصول عليها في هذه الدراسة وعلى التقييم الاقتصادي، يمكن أن نخلص إلى أن استخدام زرق الدواجن المعامل حراريا ومكبوس كمادة عضوية آمنة (ريجر) بمعدل ١٠٠ كجم /الأسبوع/ الفدان في الأشهر الثلاثة الأولى من استزراع أسماك البلطي النيلي والبوري يليها استخدام الأعلاف الصناعية التي تحتوي على البروتين الخام ٢٥٪ بمعدل ٣٪ من إجمالي وزن الجسم لأسماك البلطي النيلي و البوري في الاحواض الترابية في الأشهر الثلاثة الأخيرة يمكن أن يوصى بها لإنتاج البلطي النيلي و البوري. من وجهة النظر الإقتصادية تعتبر هذه المعاملة هي الأفضل من حيث نسبة العائد إلى إجمالي التكاليف.